

## REVIEW ABOUT COMORBIDITIES OF BEHAVIOURAL DISORDERS IN CHILDREN AND ADOLESCENTS: THE FOCUS ON ATTENTION-DEFICIT/HYPERACTIVITY DISORDER

BEATRICE GALLAI<sup>1\*</sup>, VITTORIA VALENTINI<sup>1\*</sup>, FRANCESCA BARBANERA<sup>1\*</sup>, ROSA MAROTTA<sup>2</sup>, FRANCESCO LAVANO<sup>2</sup>, SERENA MARIANNA LAVANO<sup>3</sup>, AGATA MALTESE<sup>4</sup>, GABRIELE TRIPI<sup>5,6</sup>, PALMIRA ROMANO<sup>7</sup>, MARGHERITA SALERNO<sup>8</sup>

<sup>1</sup>Department of Surgical and Biomedical Sciences, University of Perugia, Perugia, Italy - <sup>2</sup>Department of Medical and Surgery Sciences, University "Magna Graecia", Catanzaro, Italy - <sup>3</sup>Department of Health Sciences, University "Magna Graecia", Catanzaro, Italy - <sup>4</sup>Department of Psychological, Pedagogical and Educational Sciences, University of Palermo, Italy - <sup>5</sup>Department PROSAMI, University of Palermo, Italy - <sup>6</sup>Childhood Psychiatric Service for Neurodevelopmental Disorders, CH Chinon, France - <sup>7</sup>Clinic of Child and Adolescent Neuropsychiatry, Department of Mental Health and Physical and Preventive Medicine; Università degli Studi della Campania "Luigi Vanvitelli", Italy - <sup>8</sup>Sciences for Mother and Child Health Promotion, University of Palermo, Italy

\*Equal contribute for Authorship

### ABSTRACT

*Disruptive behavior disorders (DBD) present high comorbidity rate mainly for opposite-defiant disorders that are frequent among children, adolescents and adults affected by with attention deficit and hyperactivity disorder (ADHD), probably as result of common temperamental risk factors such as attention, distraction, impulsivity. ADHD tend to manifest in about 50% of individuals diagnosed as disruptive behavioral disorders.*

**Keywords:** Behavioural disorders, comorbidities, ADHD.

DOI: 10.19193/0393-6384\_2017\_2s\_186

Received November 30, 2016; Accepted May 20, 2017

### Background

Disruptive behavior disorders (DBD) present high comorbidity rate mainly for opposite-defiant disorders that are frequent among children, adolescents and adults affected by with attention deficit and hyperactivity disorder (ADHD), probably as result of common temperamental risk factors such as attention, distraction, impulsivity. ADHD tend to manifest in about 50% of individuals diagnosed as disruptive behavioral disorders<sup>(1-6)</sup>.

Actually, ADHD is included in the DSM-5 chapter of neurodevelopmental disorders and defined as a persistent pattern of inattention and/or hyperactivity-impulsivity interfering with the normal functioning or development<sup>(1-6)</sup>.

Inattention tends to manifests as inability to remain focused on a certain task, easy distraction, lack of perseverance, difficulty in paying attention to details, and organizing tasks or other daily activities. This inattention is not caused by a challenging attitude, as can be the case in provocative opposition disorder or lack of understanding<sup>(7-15)</sup>.

Hyperactivity refers to excessive motor activity in contexts and places when inappropriate, therefore the subject fails to stand still, and even when it is stopped shaking, moving hands or feet, often leaves the place in situations where he/she must be sitting (i.e. school, work), but hyperactivity can also be manifested as extreme loquacity<sup>(16-20)</sup>.

Impulsivity refers to the inability to reflect, mediate, and possibly expel behavioral responses,

taking into account the needs defined by the context. The subject does not think before acting, cannot wait for their turn in games or activities, intrudes into the games and activities of others in an intrusive way or in interrupting conversations<sup>(21-30)</sup>.

Depending on which symptom is most expressed, the DSM-5 distinguishes the ADHD in three categories: if there are both signs of inattention, hyperactivity and impulsivity, we have a combined manifestation, if there is only neglect, we talk about manifestation with predominant inattention, and finally the presence of hyperactivity and impulsiveness without any negligence outlines a manifestation with predominant hyperactivity / impulsivity<sup>(31-50)</sup>.

On the other hand, the ICD-10 defines attention deficit and hyperactivity disorder as hyperkinetic disorder (code F90), whose diagnosis requires simultaneous inattention, impulsivity and hyperactivity. Additionally, DSM-5 includes mild forms, if there are few symptoms or if they cause minor compromises in the overall functioning of the subject, serious forms if the symptoms are many or involve marked functional impairment, and moderate forms of horseback between the mild forms and those serious. Attention deficit and hyperactivity disorder begins in childhood, in fact the symptoms must occur before 12 years, without specifying an early onset due to the difficulty in establishing the exact moment in which the symptoms appear. Disturbance events must be present in multiple contexts (eg home and school, work) for more than six months, and must interfere with the normal social or work performance of the subject. The symptoms of ADHD tend to have some evolution over time: in preschool age the main manifestation is hyperactivity, while in primary school the prevalence of inattention becomes impacting<sup>(51-70)</sup>.

During adolescence, signs of hyperactivity are less common and leave more room for inner agitation, restlessness, and an inner sensation of nervousness. In adulthood tend to persist in inactivity and impulsiveness, which express themselves in making important decisions without considering the long-term consequences, in a daunting way, of accepting a job without information. Population studies indicate that ADHD is present in most cultures in about 5% of children and 2.5% in adults.

In the general population, ADHD is more common in males with an approximate ratio of 2: 1 in children and 1.6: 1 in adults. Females also tend to have males to show primarily inattention features. Among the risk factors for the development of the

ADHD plays an important role in the presence of familiarity, especially the presence of this disorder in the parents of affected people is very high<sup>(49-61)</sup>.

### *Etiology*

ADHD's inheritance is consistent even though no specific genes involved in genesis have been identified this disorder. Among the relevant causes, we can assume that the very low birth weight (<1500g) represents a doubled or tripled risk of developing this disorder, but on the other hand, most children with low birth weight will not develop ADHD. Exposure to toxic environmental factors during intrauterine life such as alcohol, cigarette smoking and childhood contact with neurotoxic substances (such as lead) or infections (encephalitis) are related to subsequent development of ADHD but It is not known whether they are related to a causal relationship. In children with this disorder can also be present a history of abuse during childhood, neglect and multiple adoptions<sup>(49-61)</sup>.

### *ADHD outcomes*

ADHD children show reduced school outcomes as a result of attentive difficulties, and will also be rejected by their companions because they are considered annoying or lazy because of their impulsive behavior and hyperactivity. Family relationships are often characterized by discord and negative interactions that, together with peer rejection, can lead to adolescence to develop comorbidity in a behavioral or mood disorder and in adulthood an antisocial personality disorder, increasing both the risk of Suicide attempts that are likely to develop substance abuse disorders and get into jail. Moreover, individuals with ADHD reach a lower level of schooling and have less personal success. In the diagnostic test, once established that symptoms of hyperactivity, impulsivity, and / or discomfort reach a level that meets the criteria for diagnosing ADHD, it is necessary to rule out that these do not fall into other diagnostic categories, often for example children with intellectual disabilities Presenting hyperkinetic behaviors, poor pulse control or easy distraction, but secondary, or cognitive-like difficulties. In children with anxiety disorders there is great difficulty in concentration and high levels of activity, even in this case secondary to an anxious background picture. A highly impulsive behavior is present in intermittent explosive disorder which is, however, rare in infancy. An increase in motor activity is also present in other neurode-

velopmental disorders such as stereotyped motion disorders and in some cases of autistic spectrum disorder, but in these cases the motions are fixed, stereotyped while in the ADHD the motor hyperactivity is typically generalized. Inattention can also be observed in children with specific learning disabilities, but in these cases it is more due to frustration and is still limited to the school context. Many children with mood disorders exhibit increased levels of activity, poor pulse control and difficulty concentrating, but in this case they are also attributed to mood disorder.

ADHD is also common as comorbid disorders, first of all the provocative opposition disorder that occurs in concomitant use in half the children with ADHD. A disorder of conduct can occur in about a quarter of children or adolescents with ADHD. The specific learning disorder commonly occurs in conjunction with ADHD, while less commonly the disorder of disruptive mood disorder, anxiety disorder, major depressive disorder, intermittent explosive disorder, obsessive-compulsive disorder, Tic disorders, and autistic spectrum disorders. Adulthood may occur in conjunction with the ADHD for substance abuse, even if in a minority of cases, antisocial personality disorder and other personality disorders. By referring to the comorbidity of disruptive behavioral disorders, often provocative opposition disturbances.

The disorder of conduct, although this seems to be more common in children with childhood recurrence. Individuals with provocative opposition disorder are also at greater risk of anxiety disorders and more depressive disorder, and this seems largely attributable to the presence of symptoms of anguish / irritability. Adolescents and adults with provocative opposition disorder also exhibit a higher rate of substance use disorders, although it is unclear whether this association is independent of comorbidity with the behavioral disorder.

How long it is common to find provocative opposition disorder and attention deficit and hyperactivity disorder, and this comorbid situation predicts the worst outcomes. Individuals with personality features associated with antisocial personality disorder often violate the fundamental rights of others or the main age-appropriate social norms, and consequently their patterns of behavior often meet the criteria for behavioral disorders. This may also coexist with one or more of specific learning disorders, anxiety disorders (depressive or bipolar), and substance-related disorders.

School outcomes, particularly in reading and other verbal skills, are often inferior to those expected by age and intelligence and can justify the additional diagnosis of specific learning disorder or communication disorder<sup>(71-100)</sup>.

## References

- 1) Muratori P, Pisano S, Milone A, Masi G. Is emotional dysregulation a risk indicator for auto-aggression behaviors in adolescents with oppositional defiant disorder? *J Affect Disord.* 2017 Jan 15; 208: 110-112. doi:10.1016/j.jad.2016.08.052.
- 2) Muratori P, Lochman JE, Lai E, Milone A, Nocentini A, Pisano S, Righini E, Masi G. Which dimension of parenting predicts the change of callous unemotional traits in children with disruptive behavior disorder? *Compr Psychiatry.* 2016 Aug; 69: 202-10. doi: 10.1016/j.comppsy.2016.06.002.
- 3) Masi G, Milone A, Manfredi A, Brovedani P, Pisano S, Muratori P. Combined pharmacotherapy-multimodal psychotherapy in children with Disruptive Behavior Disorders. *Psychiatry Res.* 2016 Apr 30; 238: 8-13. doi: 10.1016/j.psychres.2016.02.010.
- 4) Muratori P, Lochman JE, Manfredi A, Milone A, Nocentini A, Pisano S, Masi G. Callous unemotional traits in children with disruptive behavior disorder: Predictors of developmental trajectories and adolescent outcomes. *Psychiatry Res.* 2016 Feb 28; 236: 35-41. doi:10.1016/j.psychres.2016.01.003.
- 5) Chieffi S, Carotenuto M, Monda V, Valenzano A, Villano I, Precenzano F, Tafuri D, Salerno M, Filippi N, Nuccio F, Ruberto M, De Luca V, Cipolloni L, Cibelli G, Mollica MP, Iacono D, Nigro E, Monda M, Messina G and Messina A. Orexin System: The Key for a Healthy Life. *Front. Physiol.* 2017, 8:357. doi: 10.3389/fphys.2017.00357.
- 6) Pisano S, Catone G, Coppola G, Carotenuto M, Iuliano R, Tiano C, Montesanto AR, D'Esposito V, Miraglia EDG, Formisano P, Bravaccio C. Different Immune Signature in Youths Experiencing Antipsychotic-Induced Weight Gain Compared to Untreated Obese Patients. *J Child Adolesc Psychopharmacol.* 2017 Apr 28. doi: 10.1089/cap.2016.0203;
- 7) Chieffi S, Carotenuto M, Monda V, Valenzano A, Villano I, Precenzano F, Tafuri D, Salerno M, Filippi N, Nuccio F, Ruberto M, De Luca V, Cipolloni L, Cibelli G, Mollica MP, Iacono D, Nigro E, Monda M, Messina G and Messina A. Orexin System: The Key for a Healthy Life. *Front. Physiol.* 2017 8:357. doi: 10.3389/fphys.2017.00357;
- 8) Panico A, Messina G, Lupoli GA, Lupoli R, Cacciapuoti M, Moscatelli F, Esposito T, Villano I, Valenzano A, Monda V, Messina A, Precenzano F, Cibelli G, Monda M, Lupoli G. Quality of life in overweight (obese) and normal-weight women with polycystic ovary syndrome. *Patient Prefer Adherence.* 2017 Mar 2; 11: 423-429. doi: 10.2147/PPA.S119180.
- 9) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, Gallai B, Marotta R, Lavano SM, Lavano F, Roccella M. Visual-spatial training efficacy in children affected by migraine without aura: a multicenter study.

- Neuropsychiatr Dis Treat. 2017 Jan 27; 13:253-258. doi: 10.2147/NDT.S119648.
- 10) Chieffi S, Messina G, Villano I, Messina A, Esposito M, Monda V, Valenzano A, Moscatelli F, Esposito T, Carotenuto M, Viggiano A, Cibelli G, Monda M. Exercise Influence on Hippocampal Function: Possible Involvement of Orexin-A. *Front Physiol.* 2017 Feb 14; 8: 85. doi: 10.3389/fphys.2017.00085.
  - 11) Verrotti A, Casciato S, Spalice A, Carotenuto M, Striano P, Parisi P, Zamponi N, Savasta S, Rinaldi VE, D'Alonzo R, Mecarini F, Ritaccio AJ, Di Gennaro G. Coexistence of childhood absence epilepsy and benign epilepsy with centrottemporal spikes: A case series. *Eur J Paediatr Neurol.* 2017 May; 21(3): 570-575. doi: 10.1016/j.ejpn.2017.02.002.
  - 12) Villano I, Messina A, Valenzano A, Moscatelli F, Esposito T, Monda V, Esposito M, Precenzano F, Carotenuto M, Viggiano A, Chieffi S, Cibelli G, Monda M, Messina G. Basal Forebrain Cholinergic System and Orexin Neurons: Effects on Attention. *Front Behav Neurosci.* 2017 Jan 31; 11: 10. doi: 10.3389/fnbeh.2017.00010.
  - 13) Toldo I, Rattin M, Perissinotto E, De Carlo D, Bolzonella B, Nosadini M, Rossi LN, Vecchio A, Simonati A, Carotenuto M, Scalas C, Scirucchio V, Raieli V, Mazzotta G, Tozzi E, Valeriani M, Cianchetti C, Balottin U, Guidetti V, Sartori S, Battistella PA. Survey on treatments for primary headaches in 13 specialized juvenile Headache Centers: The first multicenter Italian study. *Eur J Paediatr Neurol.* 2017 May; 21(3): 507-521. doi: 10.1016/j.ejpn.2016.12.009.
  - 14) Messina A, De Fusco C, Monda V, Esposito M, Moscatelli F, Valenzano A, Carotenuto M, Viggiano E, Chieffi S, De Luca V, Cibelli G, Monda M, Messina G. Role of the Orexin System on the Hypothalamus-Pituitary-Thyroid Axis. *Front Neural Circuits.* 2016 Aug 25; 10:66. doi: 10.3389/fncir.2016.00066.
  - 15) Matricardi S, Spalice A, Salpietro V, Di Rosa G, Balistreri MC, Grosso S, Parisi P, Elia M, Striano P, Accorsi P, Cusmai R, Specchio N, Coppola G, Savasta S, Carotenuto M, Tozzi E, Ferrara P, Ruggieri M, Verrotti A. Epilepsy in the setting of full trisomy 18: A multicenter study on 18 affected children with and without structural brain abnormalities. *Am J Med Genet C Semin Med Genet.* 2016 Sep; 172(3): 288-95. doi: 10.1002/ajmg.c.31513.
  - 16) Moscatelli F, Valenzano A, Petito A, Triggiani AI, Ciliberti MAP, Luongo L, Carotenuto M, Esposito M, Messina A, Monda V, Monda M, Capranica L, Messina G, Cibelli G. Relationship between blood lactate and cortical excitability between taekwondo athletes and non-athletes after hand-grip exercise. *Somatosens Mot Res.* 2016 Jun; 33(2): 137-44. doi: 10.1080/08990220.2016.1203305.
  - 17) Carotenuto M, Esposito M, Cortese S, Laino D, Verrotti A. Children with developmental dyslexia showed greater sleep disturbances than controls including problems initiating and maintaining sleep. *Acta Paediatr.* 2016 Sep; 105(9): 1079-82. doi: 10.1111/apa.13472.
  - 18) Franzoni E, Matricardi S, Di Pisa V, Capovilla G, Romeo A, Tozzi E, Pruna D, Salerno GG, Zamponi N, Accorsi P, Giordano L, Coppola G, Cerminara C, Curatolo P, Nicita F, Spalice A, Grosso S, Pavone P, Striano P, Parisi P, Boni A, Gobbi G, Carotenuto M, Esposito M, Cottone C, Verrotti A. Refractory absence seizures: An Italian multicenter retrospective study. *Eur J Paediatr Neurol.* 2015 Nov; 19(6): 660-4. doi: 10.1016/j.ejpn.2015.07.008.
  - 19) Morandi A, Bonnefond A, Lobbens S, Carotenuto M, Del Giudice EM, Froguel P, Maffei C. A girl with incomplete Prader-Willi syndrome and negative MSPCR, found to have mosaic maternal UPD-15 at SNP array. *Am J Med Genet A.* 2015 Nov; 167A(11): 2720-6. doi: 10.1002/ajmg.a.37222.
  - 20) Pasquali D, Carotenuto M, Leporati P, Esposito M, Antinolfi L, Esposito D, Accardo G, Carella C, Chiovato L, Rotondi M. Maternal hypothyroidism and subsequent neuropsychological outcome of the progeny: a family portrait. *Endocrine.* 2015 Dec; 50(3): 797-801. doi: 10.1007/s12020-015-0564-3.
  - 21) Esposito M, Precenzano F, Sorrentino M, Avolio D, Carotenuto M. A Medical Food Formulation of Griffonia simplicifolia/Magnesium for Childhood Periodic Syndrome Therapy: An Open-Label Study on Motion Sickness. *J Med Food.* 2015 Aug; 18(8): 916-20. doi: 10.1089/jmf.2014.0113.
  - 22) Esposito M, Gallai B, Roccella M, Marotta R, Lavano F, Lavano SM, Mazzotta G, Bove D, Sorrentino M, Precenzano F, Carotenuto M. Anxiety and depression levels in prepubertal obese children: a case-control study. *Neuropsychiatr Dis Treat.* 2014 Oct 3; 10: 1897-902. doi: 10.2147/NDT.S69795.
  - 23) Verrotti A, Cusmai R, Laino D, Carotenuto M, Esposito M, Falsaperla R, Margari L, Rizzo R, Savasta S, Grosso S, Striano P, Belcastro V, Franzoni E, Curatolo P, Giordano L, Freri E, Matricardi S, Pruna D, Toldo I, Tozzi E, Lobefalo L, Operto F, Altobelli E, Chiarelli F, Spalice A. Long-term outcome of epilepsy in patients with Prader-Willi syndrome. *J Neurol.* 2015 Jan; 262(1): 116-23. doi: 10.1007/s00415-014-7542-1.
  - 24) Verrotti A, Carotenuto M, Altieri L, Parisi P, Tozzi E, Belcastro V, Esposito M, Guastaferrò N, Ciuti A, Mohn A, Chiarelli F, Agostinelli S. Migraine and obesity: metabolic parameters and response to a weight loss programme. *Pediatr Obes.* 2015 Jun; 10(3): 220-5. doi: 10.1111/ijpo.245.
  - 25) Carotenuto M, Parisi P, Esposito M, Cortese S, Elia M. Sleep alterations in children with refractory epileptic encephalopathies: a polysomnographic study. *Epilepsy Behav.* 2014 Jun; 35: 50-3. doi: 10.1016/j.yebeh.2014.03.009.
  - 26) Perillo L, Esposito M, Caprioglio A, Attanasio S, Santini AC, Carotenuto M. Orthodontic treatment need for adolescents in the Campania region: the malocclusion impact on self-concept. *Patient Prefer Adherence.* 2014 Mar 19; 8: 353-9. doi: 10.2147/PPA.S58971.
  - 27) Santamaria F, Esposito M, Montella S, Cantone E, Mollica C, De Stefano S, Mirra V, Carotenuto M. Sleep disordered breathing and airway disease in primary ciliary dyskinesia. *Respirology.* 2014 May; 19(4): 570-5. doi: 10.1111/resp.12273.
  - 28) Esposito M, Marotta R, Roccella M, Gallai B, Parisi L, Lavano SM, Carotenuto M. Pediatric neurofibromatosis 1 and parental stress: a multicenter study. *Neuropsychiatr Dis Treat.* 2014 Jan 22; 10: 141-6. doi: 10.2147/NDT.S55518.
  - 29) Esposito M, Ruberto M, Gimigliano F, Marotta R,

- Gallai B, Parisi L, Lavano SM, Roccella M, Carotenuto M. Effectiveness and safety of Nintendo Wii Fit Plus™ training in children with migraine without aura: a preliminary study. *Neuropsychiatr Dis Treat.* 2013; 9: 1803-10. doi: 10.2147/NDT.S53853.
- 30) Carotenuto M, Esposito M, Di Pasquale F, De Stefano S, Santamaria F. Psychological, cognitive and maternal stress assessment in children with primary ciliary dyskinesia. *World J Pediatr.* 2013 Nov; 9(4): 312-7. doi: 10.1007/s12519-013-0441-1.
- 31) Di Filippo T, Orlando MF, Concialdi G, La Grutta S, Lo Baido R, Epifanio MS, Esposito M, Carotenuto M, Parisi L, Roccella M. The quality of life in developing age children with celiac disease. *Minerva Pediatr.* 2013 Dec; 65(6): 599-608.
- 32) Esposito M, Parisi L, Gallai B, Marotta R, Di Dona A, Lavano SM, Roccella M, Carotenuto M. Attachment styles in children affected by migraine without aura. *Neuropsychiatr Dis Treat.* 2013; 9: 1513-9. doi: 10.2147/NDT.S52716;
- 33) Esposito M, Gimigliano F, Ruberto M, Marotta R, Gallai B, Parisi L, Lavano SM, Mazzotta G, Roccella M, Carotenuto M. Psychomotor approach in children affected by nonretentive fecal soiling (FNRFs): a new rehabilitative purpose. *Neuropsychiatr Dis Treat.* 2013; 9: 1433-41. doi: 10.2147/NDT.S51257.
- 34) Bellini B, Arruda M, Cescut A, Saulle C, Persico A, Carotenuto M, Gatta M, Nacinovich R, Piazza FP, Termine C, Tozzi E, Lucchese F, Guidetti V. Headache and comorbidity in children and adolescents. *J Headache Pain.* 2013 Sep 24; 14:79. doi:10.1186/1129-2377-14-79.
- 35) Esposito M, Roccella M, Gallai B, Parisi L, Lavano SM, Marotta R, Carotenuto M. Maternal personalità profile of children affected by migraine. *Neuropsychiatr Dis Treat.* 2013; 9: 1351-8. doi: 10.2147/NDT.S51554.
- 36) Perillo L, Esposito M, Contiello M, Lucchese A, Santini AC, Carotenuto M. Oculal traits in developmental dyslexia: a preliminary study. *Neuropsychiatr Dis Treat.* 2013; 9: 1231-7. doi: 10.2147/NDT.S49985;
- 37) Esposito M, Marotta R, Gallai B, Parisi L, Patriciello G, Lavano SM, Mazzotta G, Roccella M, Carotenuto M. Temperamental characteristics in childhood migraine without aura: a multicenter study. *Neuropsychiatr Dis Treat.* 2013; 9: 1187-92. doi: 10.2147/NDT.S50458.
- 38) Esposito M, Antinolfi L, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Mazzotta G, Precenzano F, Carotenuto M. Executive dysfunction in children affected by obstructive sleep apnea syndrome: an observational study. *Neuropsychiatr Dis Treat.* 2013; 9: 1087-94. doi: 10.2147/NDT.S47287.
- 39) Esposito M, Gallai B, Parisi L, Castaldo L, Marotta R, Lavano SM, Mazzotta G, Roccella M, Carotenuto M. Self-concept evaluation and migraine without aura in childhood. *Neuropsychiatr Dis Treat.* 2013; 9: 1061-6. doi: 10.2147/NDT.S49364.
- 40) Esposito M, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Mazzotta G, Patriciello G, Precenzano F, Carotenuto M. Visuomotor competencies and primary monosymptomatic nocturnal enuresis in prepubertal aged children. *Neuropsychiatr Dis Treat.* 2013; 9: 921-6. doi: 10.2147/NDT.S46772.
- 41) Esposito M, Parisi P, Miano S, Carotenuto M. Migraine and periodic limb movement disorders in sleep in children: a preliminary case-control study. *J Headache Pain.* 2013 Jul 1; 14: 57. doi: 10.1186/1129-2377-14-57.
- 42) Gallelli L, Avenoso T, Falcone D, Palleria C, Peltrone F, Esposito M, De Sarro G, Carotenuto M, Guidetti V. Effects of acetaminophen and ibuprofen in children with migraine receiving preventive treatment with magnesium. *Headache.* 2014 Feb; 54(2): 313-24. doi: 10.1111/head.12162.
- 43) Carotenuto M, Gimigliano F, Fiordelisi G, Ruberto M, Esposito M. Positional abnormalities during sleep in children affected by obstructive sleep apnea syndrome: the putative role of kinetic muscular chains. *Med Hypotheses.* 2013 Aug; 81(2): 306-8. doi: 10.1016/j.mehy.2013.04.023.
- 44) Esposito M, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Mazzotta G, Carotenuto M. Primary nocturnal enuresis as a risk factor for sleep disorders: an observational questionnaire-based multicenter study. *Neuropsychiatr Dis Treat.* 2013; 9: 437-43. doi: 10.2147/NDT.S43673.
- 45) Carotenuto M, Esposito M. Nutraceuticals safety and efficacy in migraine without aura in a population of children affected by neurofibromatosis type I. *Neurol Sci.* 2013 Nov; 34(11): 1905-9. doi: 10.1007/s10072-013-1403-z.
- 46) Esposito M, Carotenuto M. Intellectual disabilities and power spectra analysis during sleep: a new perspective on borderline intellectual functioning. *J Intellect Disabil Res.* 2014 May; 58(5): 421-9. doi: 10.1111/jir.12036.
- 47) Esposito M, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Gritti A, Mazzotta G, Carotenuto M. Maternal stress and childhood migraine: a new perspective on management. *Neuropsychiatr Dis Treat.* 2013; 9: 351-5. doi: 10.2147/NDT.S42818.
- 48) Esposito M, Roccella M, Parisi L, Gallai B, Carotenuto M. Hypersomnia in children affected by migraine without aura: a questionnaire-based case-control study. *Neuropsychiatr Dis Treat.* 2013; 9: 289-94. doi: 10.2147/NDT.S42182.
- 49) Parisi L, Di Filippo T, La Grutta S, Lo Baido R, Epifanio MS, Esposito M, Carotenuto M, Roccella M. Sturge-weber syndrome: a report of 14 cases. *Ment Illn.* 2013 Jun 3; 5(1): e7. doi: 10.4081/mi.2013.e7.
- 50) Carotenuto M, Gallai B, Parisi L, Roccella M, Esposito M. Acupressure therapy for insomnia in adolescents: a polysomnographic study. *Neuropsychiatr Dis Treat.* 2013; 9: 157-62. doi: 10.2147/NDT.S41892.
- 51) Esposito M, Pascotto A, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Gritti A, Mazzotta G, Carotenuto M. Can headache impair intellectual abilities in children? An observational study. *Neuropsychiatr Dis Treat.* 2012; 8: 509-13. doi:10.2147/NDT.S36863.
- 52) Carotenuto M, Esposito M, Parisi L, Gallai B, Marotta R, Pascotto A, Roccella M. Depressive symptoms and childhood sleep apnea syndrome. *Neuropsychiatr Dis Treat.* 2012; 8: 369-73. doi: 10.2147/NDT.S35974.
- 53) Esposito M, Verrotti A, Gimigliano F, Ruberto M, Agostinelli S, Scuccimarra G, Pascotto A, Carotenuto M. Motor coordination impairment and migraine in

- children: a new comorbidity? *Eur J Pediatr.* 2012 Nov;171(11): 1599-604. doi: 10.1007/s00431-012-1759-8.
- 54) Verrotti A, Agostinelli S, D'Egidio C, Di Fonzo A, Carotenuto M, Parisi P, Esposito M, Tozzi E, Belcastro V, Mohn A, Battistella PA. Impact of a weight loss program on migraine in obese adolescents. *Eur J Neurol.* 2013 Feb; 20(2): 394-7. doi: 10.1111/j.1468-1331.2012.03771.x.
- 55) Elia M, Amato C, Bottitta M, Grillo L, Calabrese G, Esposito M, Carotenuto M. An atypical patient with Cowden syndrome and PTEN gene mutation presenting with cortical malformation and focal epilepsy. *Brain Dev.* 2012 Nov; 34(10): 873-6. doi: 10.1016/j.braindev.2012.03.005.
- 56) Esposito M, Ruberto M, Pascotto A, Carotenuto M. Nutraceutical preparations in childhood migraine prophylaxis: effects on headache outcomes including disability and behaviour. *Neurol Sci.* 2012 Dec; 33(6): 1365-8. doi: 10.1007/s10072-012-1019-8.
- 57) Carotenuto M, Esposito M, D'Aniello A, Ripa CD, Precenzano F, Pascotto A, Bravaccio C, Elia M. Polysomnographic findings in Rett syndrome: a case control study. *Sleep Breath.* 2013 Mar; 17(1): 93-8. doi: 10.1007/s11325-012-0654-x. Epub 2012 Mar 7; Erratum in: *Sleep Breath.* 2013 May; 17(2): 877-8.
- 58) Guzzetta A, D'Acunto MG, Carotenuto M, Berardi N, Bancale A, Biagioni E, Boldrini A, Ghirri P, Maffei L, Cioni G. The effects of preterm infant massage on brain electrical activity. *Dev Med Child Neurol.* 2011 Sep; 53 Suppl 4:46-51. doi: 10.1111/j.1469-8749.2011.04065.x.
- 59) Carotenuto M, Esposito M, Precenzano F, Castaldo L, Roccella M. Cosleeping in childhood migraine. *Minerva Pediatr.* 2011 Apr; 63(2): 105-9.
- 60) Esposito M, Carotenuto M, Roccella M. Primary nocturnal enuresis and learning disability. *Minerva Pediatr.* 2011 Apr; 63(2): 99-104.
- 61) Esposito M, Carotenuto M. Ginkgolide B complex efficacy for brief prophylaxis of migraine in school-aged children: an open-label study. *Neurol Sci.* 2011 Feb; 32(1): 79-81. doi: 10.1007/s10072-010-0411-5.
- 62) Esposito M, Carotenuto M. Borderline intellectual functioning and sleep: the role of cyclic alternating pattern. *Neurosci Lett.* 2010 Nov 19; 485(2): 89-93. doi: 10.1016/j.neulet.2010.08.062.
- 63) Carotenuto M, Esposito M, Pascotto A. Facial patterns and primary nocturnal enuresis in children. *Sleep Breath.* 2011 May; 15(2): 221-7. doi: 10.1007/s11325-010-0388-6.
- 64) Guzzetta A, Pizzardi A, Belmonti V, Boldrini A, Carotenuto M, D'Acunto G, Ferrari F, Fiori S, Gallo C, Ghirri P, Mercuri E, Romeo D, Roversi MF, Cioni G. Hand movements at 3 months predict later emiplegia in term infants with neonatal cerebral infarction. *Dev Med Child Neurol.* 2010 Aug; 52(8): 767-72. doi: 10.1111/j.1469-8749.2009.03497.x.
- 65) Carotenuto M, Santoro N, Grandone A, Santoro E, Pascotto C, Pascotto A, Perrone L, del Giudice EM. The insulin gene variable number of tandem repeats (INS VNTR) genotype and sleep disordered breathing in childhood obesity. *J Endocrinol Invest.* 2009 Oct; 32(9): 752-5. doi: 10.3275/6398.
- 66) Elia M, Falco M, Ferri R, Spalletta A, Bottitta M, Calabrese G, Carotenuto M, Musumeci SA, Lo Giudice M, Fichera M. CDKL5 mutations in boys with severe encephalopathy and early-onset intractable epilepsy. *Neurology.* 2008 Sep 23; 71(13): 997-9. doi:10.1212/01.wnl.0000326592.37105.88.
- 67) Carotenuto M, Bruni O, Santoro N, Del Giudice EM, Perrone L, Pascotto A. Waist circumference predicts the occurrence of sleep-disordered breathing in obese children and adolescents: a questionnaire-based study. *Sleep Med.* 2006 Jun; 7(4): 357-61.
- 68) Carotenuto M, Guidetti V, Ruju F, Galli F, Tagliente FR, Pascotto A. Headache disorders as risk factors for sleep disturbances in school aged children. *J Headache Pain.* 2005 Sep; 6(4): 268-70.
- 69) Coppola G, Auricchio G, Federico R, Carotenuto M, Pascotto A. Lamotrigine versus valproic acid as first-line monotherapy in newly diagnosed typical absence seizures: an open-label, randomized, parallel-group study. *Epilepsia.* 2004 Sep; 45(9): 1049-53.
- 70) Coppola G, Licciardi F, Sciscio N, Russo F, Carotenuto M, Pascotto A. Lamotrigine as first-line drug in childhood absence epilepsy: a clinical and neurophysiological study. *Brain Dev.* 2004 Jan; 26(1): 26-9.
- 71) Capovilla G, Beccaria F, Montagnini A, Cusmai R, Franzoni E, Moscano F, Coppola G, Carotenuto M, Gobbi G, Seri S, Nabbout R, Vigeveno F. Short-term nonhormonal and nonsteroid treatment in West syndrome. *Epilepsia.* 2003 Aug; 44(8): 1085-867.
- 72) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, Vagliano C, Messina G, Di Folco A, Di Filippo T, Roccella M. Executive functioning in preschool children affected by autism spectrum disorder: a pilot study. *Acta Medica Mediterranea*, 2017, 33: 35-39; DOI: 10.19193/0393-6384\_2017\_1\_005.
- 73) Precenzano F, Lombardi P, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Della Valle I, Magliulo RM, Messina G, Roccella M. Internalizing symptoms in children affected by childhood absence epilepsy: a preliminary study. *Acta Medica Mediterranea*, 2016, 32: 1749-1753; DOI: 10.19193/0393-6384\_2016\_6\_158.
- 74) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Grappa MF, Magliulo RM, Messina G, Roccella M. Borderline intellectual functioning and parental stress: an Italian case-control study. *Acta Medica Mediterranea*, 2016, 32: 1761-1765; DOI: 10.19193/0393-6384\_2016\_6\_160.
- 75) Ruberto M, Precenzano F, Parisi L, Salerno M, Maltese A, Messina G, Roccella M. Visuomotor integration skills in children affected by obstructive sleep apnea syndrome: a case-control study. *Acta Medica Mediterranea*, 2016, 32: 1659; DOI: 10.19193/0393-6384\_2016\_5\_146.
- 76) Parisi L, Ruberto M, Precenzano F, Di Filippo T, Russotto C, Maltese A, Salerno M, Roccella M. The quality of life in children with cerebral palsy. *Acta Medica Mediterranea*, 2016, 32: 1665; DOI: 10.19193/0393-6384\_2016\_5\_147.
- 77) Epifanio, M.S., Genna, V., De Luca, C., Roccella, M., La Grutta, S. Paternal and maternal transition to parenthood. The risk of postpartum depression and parenting stress. *2015 Pediatric Reports*, 7 (2), pp. 38-44.
- 78) Parisi, L., Di Filippo, T., Roccella, M. The child with Autism Spectrum Disorders (ASDs): Behavioral and neurobiological aspects. *Acta Medica Mediterranea*,

- 2015, 31 (6), pp. 1187-1194.
- 79) Vecchio D, Salzano E, Vecchio A, Di Filippo T, Roccella, M. A case of femoral-facial syndrome in a patient with autism spectrum disorders. *Minerva Pediatrica*, 2011, 63 (4), pp. 341-344.
- 80) Parisi, L., Di Filippo, T., Roccella, M. Hypomelanosis of Ito: Neurological and psychiatric pictures in developmental age. *Minerva Pediatrica*, 2012, 64 (1), pp. 65-70.
- 81) Di Filippo, T., Parisi, L., Roccella, M. Psychological aspects in children affected by Duchenne de Boulogne muscular dystrophy. *Mental Illness*, 2012, 4 (1), pp. 21-24.
- 82) Epifanio MS, Genna V, Vitello MG, Roccella M, La Grutta S. Parenting stress and impact of illness in parents of children with coeliac disease. *Pediatr Rep*. 2013 Dec 19; 5(4): e19. doi: 10.4081/pr.2013.e19.
- 83) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Della Valle I, Visco G, Magliulo RM, Messina G, Roccella M. ADHD-like symptoms in children affected by obstructive sleep apnea syndrome: case-control study. *Acta Medica Mediterranea*, 2016, 32:1755-1759; DOI: 10.19193/0393-6384\_2016\_6\_159.
- 84) Parisi L, Salerno M, Maltese A, Tripi G, Romano P, Di Folco A, Di Filippo T, Roccella M. Anxiety levels in mothers of children affected by X-fragile syndrome. *Acta Medica Mediterranea*, 2017, 33: 495; DOI: 10.19193/0393-6384\_2017\_3\_074.
- 85) Parisi L, Salerno M, Maltese A, Tripi G, Romano P, Di Folco A, Di Filippo T, Roccella M. Autonomic regulation in autism spectrum disorders. *Acta Medica Mediterranea*, 2017, 33: 491; DOI: 10.19193/0393-6384\_2017\_3\_073.
- 86) Parisi L, Salerno M, Maltese A, Tripi G, Romano P, Di Folco A, Di Filippo T, Messina G, Roccella M. Emotional intelligence and obstructive sleep apnea syndrome in children: preliminary case-control study. *Acta Medica Mediterranea*, 2017, 33: 485; DOI: 10.19193/0393-6384\_2017\_3\_072;
- 87) Parisi L, Salerno M, Maltese A, Tripi G, Romano P, Di Folco A, Di Filippo T, Roccella M. Paternal shift-working and sleep disorders in children affected by primary nocturnal enuresis. *Acta Medica Mediterranea*, 2017, 33: 481; DOI: 10.19193/0393-6384\_2017\_3\_071;
- 88) Moscatelli F, Valenzano A, Monda V, Ruberto M, Monda G, Triggiani AI, Monda E, Chieffi S, Villano I, Parisi L, Roccella M, Messina A. Transcranial Magnetic Stimulation (TMS) application in sport medicine: A brief review. *Acta Medica Mediterranea*, 2017, 33: 423; Doi: 10.19193/0393-6384\_2017\_3\_062.
- 89) Parisi L, Faraldo Ma, Ruberto M, Salerno M, Maltese A, Di Folco A, Messina G, Di Filippo T, Roccella M. Life events and primary monosymptomatic nocturnal enuresis: a pediatric pilot study. *Acta Medica Mediterranea*, 2017, 33: 23; DOI: 10.19193/0393-6384\_2017\_1\_003.
- 90) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, Verde D, Tripi G, Romano P, Di Folco A, Di Filippo T, Messina G, Roccella M. Sleep habits in children affected by autism spectrum disorders: a preliminary case-control study. *Acta Medica Mediterranea*, 2017, 33: 405; DOI: 10.19193/0393-6384\_2017\_3\_059.
- 91) Parisi L, Fortunato MR, Salerno M, Maltese A, Di Folco A, Di Filippo T, Roccella M. Sensory perception in preschool children affected by autism spectrum disorder: A pilot study. *Acta Medica Mediterranea*, 2017, 33: 49; DOI: 10.19193/0393-6384\_2017\_1\_007;
- 92) Maltese A, Salerno M, Tripi G, Romano P, Ricciardi A, Di Folco A, Di Filippo T, Parisi L. The Angelman Syndrome: A Brief Review. *Acta Medica Mediterranea*, 2017, 33: 667; DOI: 10.19193/0393-6384\_2017\_4\_100.
- 93) Salerno M, Maltese A, Tripi G, Romano P, Di Folco A, Di Filippo T. Separation anxiety in pediatric migraine without aura: A Pilot Study. *Acta Medica Mediterranea*, 2017, 33: 621; DOI: 10.19193/0393-6384\_2017\_4\_092.
- 94) Maltese A, Salerno M, Tripi G, Romano P, Ricciardi A, Sessa G, Di Folco A, Di Filippo T, Parisi L. Rehabilitative treatment proposals in pediatric non-verbal syndrome. *Acta Medica Mediterranea*, 2017, 33: 675; DOI: 10.19193/0393-6384\_2017\_4\_101.
- 95) Monda V, Nigro E, Ruberto M, Monda G, Valenzano A, Triggiani Ai, Moscatelli F, Monda E, Villano I, Roccella M, Parisi L, Messina A. Synergism or competition between zinc and chromium dietary levels on insulin action mechanism. A method to investigate. *Acta Medica Mediterranea*, 2017, 33: 581; DOI: 10.19193/0393-6384\_2017\_4\_085.
- 96) Messina A, Monda V, Avola R, Moscatelli F, Valenzano AA, Villano I, Ruberto M, Monda E, La Marra M, Tafuri D, Chieffi S, Cibelli G, Monda M, Messina G. Role of the orexin system on arousal, attention, feeding behaviour and sleep disorders. *Acta Medica Mediterranea*, 2017, 33: 645; DOI: 10.19193/0393-6384\_2017\_4\_096.
- 97) Messina A, Monda V, Nigro E, Valenzano AA; Villano I, Ruberto M, Monda G, Ascione A, Chieffi S, Cibelli G, Messina G, Monda M. AN Allied health: the pasta. *Acta Medica Mediterranea*, 2017, 33: 641; DOI: 10.19193/0393-6384\_2017\_4\_095.
- 98) Messina A, Russo G, Monda V, Valenzano A, Villano I, Ascione A, Moscatelli F, Crescenzo R, Catizzone Ar, Panico A, Fulgione E, Piombino L, Dorato D, Cavaliere G, Trinchese G, Cibelli G, Bartoletti E, Messina G. Effect of radiofrequency on sympathetic nervous system functioning. *Acta Medica Mediterranea*, 2017, 33: 833; DOI: 10.19193/0393-6384\_2017\_5\_124.
- 99) Chisari G, Rampello L, Chisari EM, Catania VE, Greco C, Stagni E, Chisari CG. Microbiology synergism between tear substitutes and symbiotic treatment of patients with irritable bowel syndrome. *Acta Medica Mediterranea*, 2016 32 (4), pp. 865-870. Cited 1 time. DOI: 10.19193/0393-6384\_2016\_4\_102.
- 100) Wen S, Jin Y, Wang L. Comparison study of integrin B2 gene expressions between acute myocardial infarction and stable angina pectoris patients. *Acta Medica Mediterranea*, 2016 32 (2), pp. 303-312. DOI: 10.19193/0393-6384\_2016\_2\_45.

*Corresponding author*

MARGHERITA SALERNO, MD

Sciences for Mother and Child Health Promotion

University of Palermo

(Italy)