

THE SYNACTIVE THEORY OF DEVELOPMENT: THE KEYWORD FOR NEURODEVELOPMENTAL DISORDERS

AGATA MALTESE^{1*}, BEATRICE GALLAI^{2*}, ROSA MAROTTA^{3*}, FRANCESCO LAVANO³, SERENA MARIANNA LAVANO³, GABRIELE TRIPI^{4,5}, PALMIRA ROMANO⁶, LUCREZIA D'ORO⁶, MARGHERITA SALERNO⁷

¹Department of Psychological, Pedagogical and Educational Sciences, University of Palermo, Italy - ²Department of Surgical and Biomedical Sciences, University of Perugia, Perugia, Italy - ³Department of Medical and Surgery Sciences, University "Magna Graecia", Catanzaro, Italy - ⁴Department PROSAMI, University of Palermo, Italy - ⁵Childhood Psychiatric Service for Neurodevelopmental Disorders, CH Chinon, France - ⁶Clinic of Child and Adolescent Neuropsychiatry, Department of Mental Health and Physical and Preventive Medicine; Università degli Studi della Campania "Luigi Vanvitelli", Italy - ⁷Sciences for Mother and Child Health Promotion, University of Palermo, Italy

**Equal contribution*

ABSTRACT

The synactive theory of development may be considered the keyword for neurodevelopmental disorders, considering that each one presents constantly autonomic troubles such as sleep disorders, feeding problems.

Keywords: *Synactive Theory of Development, autonomic nervous system, sleep.*

DOI: 10.19193/0393-6384_2017_2s_194

Received December 30, 2016; Accepted June 20, 2017

Introduction

The nervous system is distinct in the central nervous system (CNS), consisting of the brainstem, spinal cord and peripheral nervous system (PNS), which includes all nerve structures located outside the CNS. PNS is composed by an afferent component for sensitive information, and an efferent component for peripheral organs control. Moreover, efferent system is composed by somatic component (for skeletal muscles contraction regulation and control) and visceral or autonomic component identified as neurovegetative system (for heart and respiratory frequencies, digestive, urinary, and reproductive functions regulation). In neurodevelopmental disorders constantly⁽¹⁻⁶⁾.

The Synactive Theory

The synactive development (SDT) model was developed by Heidelise Als, American neuropsychologist in Boston, providing a different access to brain research through child's behavior observation. In this picture, behavior is the first way for preterm infants and neonate may communicate and by simple observing, it is possible to put in place adequate care paths.

Assistance process is understood as a co-regulation and collaboration action to enable and maintain the homeostasis of the various subsystems. SDT enhances the way the child appears to handle the experiences from the environment rather than the individual abilities⁽¹⁻⁶⁾.

SDT identifies development as an interactive and hierarchical process including five subsystems:

- 1) neurovegetative/autonomic system,
- 2) motor system,
- 3) behavioral system,
- 4) attention-to-interaction system,
- 5) self-regulation system (Figure 1).




Figure 1: The synactive theory development scheme.

These systems develop in the embryo, in the fetus and in the newborn according to a well-established sequence, according to independent development. Nevertheless, the 5 systems are constantly influenced, although interdependent and their development takes place interactively with the environment. The good functioning and stability of each system facilitates the maturation of the next one as well as the instability and disorganization of a system adversely affect the functioning and maturation of others.

In order of development we have:

- Nervous system: observable through vital functions, therefore with respiratory and cardiac frequency, thermoregulation, cutaneous color, visceral functions such as excretory, regurgitation, hiccups, vomiting and motor functions such as tremors, startles and clones;

- Motor system evaluated by observing the different postures taken by the newborn (hands to the mouth, hands to the face, bracing, grasping, sucking, cuddling), the quality and variability of the movements of the limbs and trunk and observing all strategies for self-regulation. Arm /trunk movements (quality, quantity);

- Behavioral states: the status of states of con-

sciousness, the state of sleep, wakefulness and agitation, the mode of transition from one state to the other, and the presence and duration of the individual states are observed. The states of consciousness are: deep sleep (quiet); Light sleep (active - REM); Drowsiness (sleeping); Calm / active, that is, the state necessary to interact with the child, characterized by reduced motor activity, opening and enlarging the eyes, attentive look in response to auditory / visual stimuli, regular respiratory rate; Shaken Vigil; Cry.

- Attention-Interaction System: Assess the ability of the child to maintain a state of vigilant alert that can be observed through the ability to relate to the environment and caregiver;

- Self-regulation system: evaluable, after 36 weeks gestational age, through the strategies the body uses to find a stable balance and the organization of the interaction between the subsystems. If the infant can not find a stable interaction between the subsystems, the quality and type of environmental facilitation must be assessed to help the child reach this organization

In neuromotor development these subsystems mature according to a compulsory sequence, divided into three phases, during each one, there are precise characteristics at each state level (1-6).

The Stabilization Phase (24 to 29 weeks gestational age):

- Neurovegetative/autonomic System is unstable and breathing and feeding are not autonomous,

- Motor System is unstable and the postural control is poor;

- Behavioral System is not evaluated because there are cycles of alternation of movement and quiet and an alternation of sleep/wake;

- Attention-Interaction System → the Interactive skills are developed during the fetal life. Fetus has a perception and reaction to stimuli through tactile, proprioceptive, painful, auditory, vestibular sensitivity. After born, the interaction takes place thanks to the body language and thanks to the different visual, olfactory, tasting, and thermal devices.

- Self-regulation system is still very poor

The Organization Phase (30 to 35 weeks gestational age):

- Neurovegetative System is more stable, respiration often becomes autonomous, as feeding, linked to the coordination of suction, swallowing and breathing functions •

- Motor System is more stable with postural

control that begins to mature with the acquisition of antigravity skills

- Behavioral System → now we can recognize the different states, there begins to be an alternation between sleep and wake, between tears and consolation, between the state of quiet and the movement and, thanks to the acquisition of visual skills, also appears the alert state •

- Attentiveness-Interaction System. The attentive and alertness skills appear more stable and the child is capable of short social interactions, succeeding in responding to a stimulus at a time •

- Self-regulation system begins to have a hint of maturation but there is a need for facilitations

The Integration phase (36-40 weeks gestational age):

- Neurovegetative/autonomic system is still Stable The respiratory system and the feeding are autonomous;

- Motor System is even more stable with a postural control that is acquired, as well as the most antigravitational skills that are observable in the various postures, prong, supine

- Behavioral System has a precise definition with a well-defined cyclical organization with the alternation of different sleep/wake states, crying/consolation, rest/movement

- Attention-Interaction System. The social interactions are more stable and greater propositiveness can be found • self-regulation system

- Self-regulation capacity increases and there is a need for external facilitations⁽⁷⁻⁵⁷⁾.

The goals of SDT are many; firstly, we want to propose reading the behavior of the infant through signs of self-regulation or, conversely, of stress of the know to provide a tool for a more detailed and sophisticated observation of the infant's behavior, to fully understand the development, and finally to plan customized care measures tailored to that individual's child in order to promote their development

Through the facilitation of the organization, the maturation and integration of the individual subsystems. Stress caused by care maneuvers, invasiveness of environmental stimuli and medical interventions causes a destabilization of the entire premature organism. Neuro-vascular, motor, behavioral, attention-interaction and self-regulation skills can be observed to specifically identify stress tolerance thresholds, level of organization, evolving capabilities, self-regulation and self-differentiation. The baby's world is full of new experiences that the child learns to handle quickly.

The infant seems to experience the world significantly, learning to quickly adjust his own behavior, aiming to express his intentions and requests for support. The infant reaches these behavioral goals through self-regulation support and the positive integration and interaction of the five subsystems. Systems are interdependent and the behaviors that express them are significant and communicative. The behaviors of a newborn, but in general of the child, indicate the level of self-regulation and the contribution of each behavioral system to this regulatory process. Self-regulation is always determined in the context of an environmental event, which can also be a physical therapy intervention.

Environmental events call upon the infant and challenge her ability to maintain a state of good behavioral organization. Adequate self-directed behaviors indicate a child's state of well-being and are observed when self-regulation is able to support the social and environmental demands that are proposed to him/her. Loss of self-regulation and disorganization manifestations are a vulnerability to the child's development, or what the child has difficulty in doing under specific conditions. While stress behaviors indicate a state of tiredness or disorganization and have been observed when self-regulating thresholds in the newborn are overcome by the situation being proposed.

A bimodal approach to understanding neuro-behavioral functioning can, however, only partly explain the behaviors that manifest themselves as transitions from a self-regulating state to one of a loss of self-regulation. Stress or elusive behavior requires a lot of attention from the 'Observer, as at the moment when the child's sensory thresholds are reached, it is necessary to offer support to the baby to minimize or prevent the complete loss of self-regulation⁽⁵⁸⁻⁸⁵⁾.

This model may be applied in pediatric management of neurodevelopmental disorders, in order to improve the quality care.

References

- 1) Als, H (1986). "A synactive model of neonatal behavioral organization: Framework for the assessment and support of the neurobehavioral development of the premature infant and his parents in the environment of the neonatal intensive care unit. In JK Sweeney (ed.), *The high risk neonate: Developmental therapy perspectives*". *Physical and Occupational Therapy in Pediatrics*. 6: 3-53;

- 2) Martini R, Cramm H, Egan M, Sikora L. Scoping Review of Self-Regulation: What Are Occupational Therapists Talking About? *Am J Occup Ther.* 2016 Nov/Dec;70(6):7006290010p1-7006290010p15.
- 3) Bembich S, Fiani G, Strajn T, Sanesi C, Demarini S, Sanson G. Longitudinal Responses to Weighing and Bathing Procedures in Preterm Infants. *J Perinat Neonatal Nurs.* 2017 Jan/Mar; 31(1): 67-74. Doi:10.1097/JPN.0000000000000228.
- 4) Roan C, Bell A. Occupational Therapy in the Neonatal Intensive Care Unit for a Neonate with Perinatal Stroke: A Case Report. *Phys Occup Ther Pediatr.* 2017 Aug; 37(3): 283-291. doi: 10.1080/01942638.2016.1185503.
- 5) Gibbins S, Hoath SB, Coughlin M, Gibbins A, Frank L. The universe of developmental care: a new conceptual model for application in the neonatal intensive care unit. *Adv Neonatal Care.* 2008 Jun; 8(3): 141-7. doi: 10.1097/01.ANC.0000324337.01970.76.
- 6) Pickler RH. A Model of Feeding Readiness for Preterm Infants. *Neonatal Intensive Care.* 2004; 17(4): 31-36;.
- 7) Messina, G., Palmieri, F., Monda, V., Messina, A., Dalia, C., Viggiano, A., Tafuri, D., Messina, A., Moscatelli, F., Valenzano, A., Cibelli, G., Chieffi, S., Monda, M. Exercise Causes Muscle GLUT4 Translocation in an Insulin-Independent Manner. *Biol Med.* 2015;s3. doi:10.4172/0974-8369.1000s3007.
- 8) Valenzano A, Moscatelli F, Triggiani AI, Capranica L, De Ioannon G, Piacentini MF, Mignardi S, Messina G, Villani S, Cibelli G. Heart-Rate Changes After an Ultraendurance Swim From Italy to Albania: A Case Report. *Int J Sports Physiol Perform.* 2016; 11(3): 407-9. doi:10.1123/ijsp.2015-0035.
- 9) Moscatelli, F., Messina, G., Valenzano, A., Petito, A., Triggiani, A.I., Ciliberti, M.A.P., Monda, V., Messina, A., Tafuri, D., Capranica, L., Cibelli, G., Monda, M. Relationship between RPE and Blood Lactate after Fatiguing Handgrip Exercise in Taekwondo and Sedentary Subjects. *Biol Med.* 2015. doi:http://dx.doi.org/10.4172/0974-8369.1000s3008;
- 10) Rinaldi B, Guida F, Furiano A, Donniacuo M, Luongo L, Gritti G, Urbanek K, Messina G, Maione S, Rossi F, de Novellis V. Effect of Prolonged Moderate Exercise on the Changes of Nonneuronal Cells in Early Myocardial Infarction. *Neural Plast.* 2015; 2015:265967. doi:10.1155/2015/265967;
- 11) Messina, G., Viggiano, A., Tafuri, D., Palmieri, F., De Blasio, S., Messina, A., De Luca, A., Chieffi, S., Monda, M. Role of orexin in obese patients in the intensive care unit. *J Anesth Clin Res.* 2014;5(3). doi:10.4172/2155-6148.1000395;
- 12) Messina, G., Zannella, C., Monda, V., Dato, A., Liccardo, D., De Blasio, S., Valenzano, A., Moscatelli, F., Messina, A., Cibelli, G., Monda, M. The Beneficial Effects of Coffee in Human Nutrition. 2015; 7(4). doi:10.4172/0974-8369.1000.
- 13) Messina A, De Fusco C, Monda V, Esposito M, Moscatelli F, Valenzano A, Carotenuto M, Viggiano E, Chieffi S, De Luca V, Cibelli G, Monda M, Messina G. Role of the Orexin System on the Hypothalamus-Pituitary-Thyroid Axis. *Front Neural Circuits.* 2016; 10: 66. doi:10.3389/fncir.2016.00066.
- 14) Mazzeo, F., Motti, M.L., Messina, G., Monda, V., Ascione, A., Tafuri, D., Palmieri, F., Messina, A., Monda, M. Use of nutritional supplements among south Italian students of physical training and sport university. *Curr Top Toxicol.* 2013 9: 21-26.
- 15) Chieffi, S., Iavarone, A., La Marra, M., Messina, G., Dalia, C., Viggiano, A., De Luca, V., Monda, M. Vulnerability to Distraction in Schizophrenia. *J PSYCHIATRY.* 2015; 18(2). doi:10.4172/1994-8220.1000228.
- 16) Esposito, T., Viggiano, An., Viggiano, Al., Viggiano, E., Giovane, A., Varriale, B., Messina, G., De Luca, V., Monda, M. ICV injection of orexin A induces synthesis of total RNA and mRNA encoding preorexin in various cerebral regions of the rat. *J Therm Biol.* 2006; 31(7): 527-532. doi:10.1016/j.jtherbio.2006.07.002.
- 17) Messina G, Di Bernardo G, Viggiano A, De Luca V, Monda V, Messina A, Chieffi S, Galderisi U, Monda M. Exercise increases the level of plasma orexin A in humans. *J Basic Clin Physiol Pharmacol.* 2016; 27(6): 611-616. doi:10.1515/jbcpp-2015-0133.
- 18) Viggiano E, Monda V, Messina A, Moscatelli F, Valenzano A, Tafuri D, Cibelli G, De Luca B, Messina G, Monda M. Cortical spreading depression produces a neuroprotective effect activating mitochondrial uncoupling protein-5. *Neuropsychiatr Dis Treat.* 2016;12:1705-10. doi:10.2147/NDT.S107074;
- 19) Moscatelli F, Messina G, Valenzano A, Monda V, Viggiano A, Messina A, Petito A, Triggiani AI, Ciliberti MA, Monda M, Capranica L, Cibelli G. Functional Assessment of Corticospinal System Excitability in Karate Athletes. *PLoS One.* 2016; 11(5): e0155998. doi:10.1371/journal.pone.0155998;
- 20) Messina, G., Valenzano, A., Moscatelli, F., Triggiani, A.I., Capranica, L., Messina, A., Piombino, L., Tafuri, D., Cibelli, G., Monda, M. Effects of Emotional Stress on Neuroendocrine and Autonomic Functions in Skydiving. *J PSYCHIATRY.* 2015; 18(4). doi:10.4172/2378-5756.1000280.
- 21) Esposito M, Carotenuto M. Intellectual disabilities and power spectra analysis during sleep: a new perspective on borderline intellectual functioning. *J Intellect Disabil Res.* 2014 May; 58(5): 421-9. doi: 10.1111/jir.12036;
- 22) Coppola G, Licciardi F, Sciscio N, Russo F, Carotenuto M, Pascotto A. Lamotrigine as first-line drug in childhood absence epilepsy: a clinical and neurophysiological study. *Brain Dev.* 2004 Jan; 26(1): 26-9;
- 23) Carotenuto M, Esposito M. Nutraceuticals safety and efficacy in migraine without aura in a population of children affected by neurofibromatosis type I. *Neurol Sci.* 2013 Nov; 34(11): 1905-9. doi: 10.1007/s10072-013-1403-z.
- 24) Carotenuto M, Gimigliano F, Fiordelisi G, Ruberto M, Esposito M. Positional abnormalities during sleep in children affected by obstructive sleep apnea syndrome: the putative role of kinetic muscular chains. *Med Hypotheses.* 2013 Aug; 81(2): 306-8. doi: 10.1016/j.mehy.2013.04.023;
- 25) Perillo L, Esposito M, Caprioglio A, Attanasio S, Santini AC, Carotenuto M. Orthodontic treatment need for adolescents in the Campania region: the malocclusion impact on self-concept. *Patient Prefer Adherence.* 2014 Mar 19; 8: 353-9. doi: 10.2147/PPA.S58971.
- 26) Carotenuto M, Esposito M, Pascotto A. Migraine and enuresis in children: An unusual correlation? *Med*

- Hypotheses. 2010 Jul; 75(1): 120-2. doi: 10.1016/j.mehy.2010.02.004;
- 27) Esposito M, Gallai B, Roccella M, Marotta R, Lavano F, Lavano SM, Mazzotta G, Bove D, Sorrentino M, Precenzano F, Carotenuto M. Anxiety and depression levels in prepubertal obese children: a case-control study. *Neuropsychiatr Dis Treat*. 2014 Oct 3;10:1897-902. doi: 10.2147/NDT.S69795.
- 28) Esposito M, Parisi L, Gallai B, Marotta R, Di Dona A, Lavano SM, Roccella M, Carotenuto M. Attachment styles in children affected by migraine without aura. *Neuropsychiatr Dis Treat*. 2013;9:1513-9. doi: 10.2147/NDT.S52716.
- 29) Messina A, Monda V, Avola R, Moscatelli F, Valenzano AA, Villano I, Ruberto M, Monda E, La Marra M, Tafuri D, Chieffi S, Cibelli G, Monda M, Messina G. Role of the orexin system on arousal, attention, feeding behaviour and sleep disorders. *Acta Medica Mediterranea*, 2017, 33: 645; DOI: 10.19193/0393-6384_2017_4_096.
- 30) Carotenuto M, Esposito M, Cortese S, Laino D, Verrotti A. Children with developmental dyslexia showed greater sleep disturbances than controls, including problems initiating and maintaining sleep. *Acta Paediatr*. 2016 Sep;105(9):1079-82. doi: 10.1111/apa.13472.
- 31) Santamaria F, Esposito M, Montella S, Cantone E, Mollica C, De Stefano S, Mirra V, Carotenuto M. Sleep disordered breathing and airway disease in primary ciliary dyskinesia. *Respirology*. 2014 May; 19(4): 570-5. doi: 10.1111/resp.12273.
- 32) Esposito M, Gallai B, Parisi L, Castaldo L, Marotta R, Lavano SM, Mazzotta G, Roccella M, Carotenuto M. Self-concept evaluation and migraine without aura in childhood. *Neuropsychiatr Dis Treat*. 2013; 9: 1061-6. doi: 10.2147/NDT.S49364.
- 33) Esposito M, Roccella M, Gallai B, Parisi L, Lavano SM, Marotta R, Carotenuto M. Maternal personality profile of children affected by migraine. *Neuropsychiatr Dis Treat*. 2013; 9: 1351-8. doi: 10.2147/NDT.S51554.
- 34) Esposito M, Marotta R, Gallai B, Parisi L, Patriciello G, Lavano SM, Mazzotta G, Roccella M, Carotenuto M. Temperamental characteristics in childhood migraine without aura: a multicenter study. *Neuropsychiatr Dis Treat*. 2013; 9: 1187-92. doi: 10.2147/NDT.S50458.
- 35) Esposito M, Precenzano F, Sorrentino M, Avolio D, Carotenuto M. A Medical Food Formulation of Griffonia simplicifolia/Magnesium for Childhood Periodic Syndrome Therapy: An Open-Label Study on Motion Sickness. *J Med Food*. 2015 Aug;18(8):916-20. doi: 10.1089/jmf.2014.0113.
- 36) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, Vagliano C, Messina G, Di Folco A, Di Filippo T, Roccella M. Executive functioning in preschool children affected by autism spectrum disorder: a pilot study. *Acta Medica Mediterranea*, 2017, 33: 35-39; DOI: 10.19193/0393-6384_2017_1_005.
- 37) Precenzano F, Lombardi P, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Della Valle I, Magliulo RM, Messina G, Roccella M. Internalizing symptoms in children affected by childhood absence epilepsy: a preliminary study. *Acta Medica Mediterranea*, 2016, 32: 1749-1753; DOI: 10.19193/0393-6384_2016_6_158.
- 38) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Grappa MF, Magliulo RM, Messina G, Roccella M. Borderline intellectual functioning and parental stress: an Italian case-control study. *Acta Medica Mediterranea*, 2016, 32: 1761-1765; DOI: 10.19193/0393-6384_2016_6_160.
- 39) Ruberto M, Precenzano F, Parisi L, Salerno M, Maltese A, Messina G, Roccella M. Visuomotor integration skills in children affected by obstructive sleep apnea syndrome: a case-control study. *Acta Medica Mediterranea*, 2016, 32: 1659; DOI: 10.19193/0393-6384_2016_5_146;
- 40) Parisi L, Ruberto M, Precenzano F, Di Filippo T, Russotto C, Maltese A, Salerno M, Roccella M. The quality of life in children with cerebral palsy. *Acta Medica Mediterranea*, 2016, 32: 1665; DOI: 10.19193/0393-6384_2016_5_147.
- 41) Epifanio, M.S., Genna, V., De Luca, C., Roccella, M., La Grutta, S. Paternal and maternal transition to parenthood. The risk of postpartum depression and parenting stress. 2015 *Pediatric Reports*, 7 (2), pp. 38-44.
- 42) Parisi, L., Di Filippo, T., Roccella, M. The child with Autism Spectrum Disorders (ASDs): Behavioral and neurobiological aspects. *Acta Medica Mediterranea*, 2015, 31 (6), pp. 1187-1194.
- 43) Vecchio D, Salzano E, Vecchio A, Di Filippo T, Roccella, M. A case of femoral-facial syndrome in a patient with autism spectrum disorders. *Minerva Pediatrica*, 2011, 63 (4), pp. 341-344.
- 44) Parisi, L., Di Filippo, T., Roccella, M. Hypomelanosis of Ito: Neurological and psychiatric pictures in developmental age. *Minerva Pediatrica*, 2012, 64 (1), pp. 65-70.
- 45) Di Filippo, T., Parisi, L., Roccella, M. Psychological aspects in children affected by Duchenne de Boulogne muscular dystrophy. *Mental Illness*, 2012, 4 (1), pp. 21-24.
- 46) Epifanio MS, Genna V, Vitello MG, Roccella M, La Grutta S. Parenting stress and impact of illness in parents of children with coeliac disease. *Pediatr Rep*. 2013 Dec 19; 5(4): e19. doi: 10.4081/pr.2013.e19.
- 47) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Della Valle I, Visco G, Magliulo RM, Messina G, Roccella M. ADHD-like symptoms in children affected by obstructive sleep apnea syndrome: case-control study. *Acta Medica Mediterranea*, 2016, 32:1755-1759; DOI: 10.19193/0393-6384_2016_6_159.
- 48) Parisi L, Salerno M, Maltese A, Tripi G, Romano P, Di Folco A, Di Filippo T, Roccella M. Anxiety levels in mothers of children affected by X-fragile syndrome. *Acta Medica Mediterranea*, 2017, 33: 495; DOI: 10.19193/0393-6384_2017_3_074.
- 49) Parisi L, Salerno M, Maltese A, Tripi G, Romano P, Di Folco A, Di Filippo T, Roccella M. Autonomic regulation in autism spectrum disorders. *Acta Medica Mediterranea*, 2017, 33: 491; DOI: 10.19193/0393-6384_2017_3_073.
- 50) Parisi L, Salerno M, Maltese A, Tripi G, Romano P, Di Folco A, Di Filippo T, Messina G, Roccella M. Emotional intelligence and obstructive sleep apnea syndrome in children: preliminary case-control study. *Acta*

- Medica Mediterranea, 2017, 33: 485; DOI: 10.19193/0393-6384_2017_3_072.
- 51) Parisi L, Salerno M, Maltese A, Tripi G, Romano P, Di Folco A, Di Filippo T, Roccella M. Paternal shift-working and sleep disorders in children affected by primary nocturnal enuresis. *Acta Medica Mediterranea*, 2017, 33: 481; DOI: 10.19193/0393-6384_2017_3_071.
- 52) Moscatelli F, Valenzano A, Monda V, Ruberto M, Monda G, Triggiani AI, Monda E, Chieffi S, Villano I, Parisi L, Roccella M, Messina A. Transcranial Magnetic Stimulation (TMS) application in sport medicine: A brief review. *Acta Medica Mediterranea*, 2017, 33: 423; Doi: 10.19193/0393-6384_2017_3_062.
- 53) Parisi L, Faraldo Ma, Ruberto M, Salerno M, Maltese A, Di Folco A, Messina G, Di Filippo T, Roccella M. Life events and primary monosymptomatic nocturnal enuresis: a pediatric pilot study. *Acta Medica Mediterranea*, 2017, 33: 23; DOI: 10.19193/0393-6384_2017_1_003.
- 54) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, Verde D, Tripi G, Romano P, Di Folco A, Di Filippo T, Messina G, Roccella M. Sleep habits in children affected by autism spectrum disorders: a preliminary case-control study. *Acta Medica Mediterranea*, 2017, 33: 405; DOI: 10.19193/0393-6384_2017_3_059.
- 55) Parisi L, Fortunato MR, Salerno M, Maltese A, Di Folco A, Di Filippo T, Roccella M. Sensory perception in preschool children affected by autism spectrum disorder: A pilot study. *Acta Medica Mediterranea*, 2017, 33: 49; DOI: 10.19193/0393-6384_2017_1_007.
- 56) Maltese A, Salerno M, Tripi G, Romano P, Ricciardi A, Di Folco A, Di Filippo T, Parisi L. The Angelman Syndrome: A Brief Review. *Acta Medica Mediterranea*, 2017, 33: 667; DOI: 10.19193/0393-6384_2017_4_100.
- 57) Salerno M, Maltese A, Tripi G, Romano P, Di Folco A, Di Filippo T. Separaxion anxiety in pediatric migraine without aura: A Pilot Study. *Acta Medica Mediterranea*, 2017, 33: 621; DOI: 10.19193/0393-6384_2017_4_092.
- 58) Maltese A, Salerno M, Tripi G, Romano P, Ricciardi A, Sessa G, Di Folco A, Di Filippo T, Parisi L. Rehabilitative treatment proposals in pediatric non-verbal syndrome. *Acta Medica Mediterranea*, 2017, 33: 675; DOI: 10.19193/0393-6384_2017_4_101.
- 59) Monda V, Nigro E, Ruberto M, Monda G, Valenzano A, Triggiani Ai, Moscatelli F, Monda E, Villano I, Roccella M, Parisi L, Messina A. Synergism or competition between zinc and chromium dietary levels on insulin action mechanism. A method to investigate. *Acta Medica Mediterranea*, 2017, 33: 581; DOI: 10.19193/0393-6384_2017_4_085.
- 60) Messina A, Monda V, Avola R, Moscatelli F, Valenzano AA, Villano I, Ruberto M, Monda E, La Marra M, Tafuri D, Chieffi S, Cibelli G, Monda M, Messina G. Role of the orexin system on arousal, attention, feeding behaviour and sleep disorders. *Acta Medica Mediterranea*, 2017, 33: 645; DOI: 10.19193/0393-6384_2017_4_096.
- 61) Messina A, Monda V, Nigro E, Valenzano AA, Villano I, Ruberto M, Monda G, Ascione A, Chieffi S, Cibelli G, Messina G, Monda M. AN Allied health: the pasta. *Acta Medica Mediterranea*, 2017, 33: 641; DOI: 10.19193/0393-6384_2017_4_095.
- 62) Messina A, Russo G, Monda V, Valenzano A, Villano I, Ascione A, Moscatelli F, Crescenzo R, Catizzone Ar, Panico A, Fulgione E, Piombino L, Dorato D, Cavaliere G, Trinchese G, Cibelli G, Bartoletti E, Messina G. Effect of radiofrequency on sympathetic nervous system functioning. *Acta Medica Mediterranea*, 2017, 33: 833; DOI: 10.19193/0393-6384_2017_5_124.
- 63) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, Gallai B, Marotta R, Lavano SM, Lavano F, Roccella M. Visual-spatial training efficacy in children affected by migraine without aura: a multicenter study. *Neuropsychiatr Dis Treat*. 2017 Jan 27; 13: 253-258. doi: 10.2147/NDT.S119648.
- 64) Maltese, A., Salerno, M., Romano, P., Ricciardi, A., Di Filippo, T., Tripi, G. Ketogenic diet as antiepileptic therapy: Administration and formulas. *Acta Medica Mediterranea*, 2017 33 (5), pp. 737-744. DOI: 10.19193/0393-6384_2017_5_108.
- 65) Maltese, A., Salerno, M., Romano, P., Ricciardi, A., Di Filippo, T., Tripi, G. Ketogenic diet as antiepileptic therapy: Neurotransmission effects. *Acta Medica Mediterranea*, 2017 33 (5), pp. 777-783. DOI: 10.19193/0393-6384_2017_5_114.
- 66) Maltese, A., Salerno, M., Romano, P., Ricciardi, A., di Filippo, T., Tripi, G. Ketogenic diet as antiepileptic therapy: Historical perspective. *Acta Medica Mediterranea*, 2017 33 (5), pp. 769-775.
- 67) Maltese, A., Salerno, M., Tripi, G., Romano, P., Ricciardi, A., di Folco, A., di Filippo, T., Parisi, L. Internalizing problems are related to sleep patterns disordered in children affected by primary headache. *Acta Medica Mediterranea*, 2017 33 (5), pp. 729-735. DOI: 10.19193/0393-6384_2017_5_107.
- 68) Salerno, M., Maltese, A., Gabrieletripi, Romano, P., Di Folco, A., Di Filippo, T., Parisi, L. Separaxion anxiety in pediatric migraine without aura: A pilot study. *Acta Medica Mediterranea*, 2017 33 (4), pp. 621-625. DOI: 10.19193/0393-6384_2017_4_092.
- 69) Maltese, A., Salerno, M., Tripi, G., Romano, P., Ricciardi, A., Sessa, G., Di Folco, A., Di Filippo, T., Parisi, L. Rehabilitative treatment proposals in pediatric non-verbal syndrome. *Acta Medica Mediterranea*, 2017 33 (4), p. 675. DOI: 10.19193/0393-6384_2017_4_101.
- 70) Giardina, S., Montinari, M.R. The incurable wounds of the soul: Suicide as a tragic self-affirmation in art and literature (2016) *Acta Medica Mediterranea*, 32 (6), art. no. 1739. DOI: 10.19193/0393-6384_2016_6_157.
- 71) Bosnak, V., Karaoglan, I., Namiduru, M. Vertebral osteomyelitis: Seven years experience of 85 cases (2016) *Acta Medica Mediterranea*, 32 (4), pp. 997-1002. DOI: 10.19193/0393-6384_2016_4_122.
- 72) Özçelik, K.Ç., Ulucanlar, H., Turanlı, S., Zihni, I., Karaköse, O., Pülüt, H., Çalta, A.F., Eken, H. Clinical significance of the multicentric papillary thyroid carcinoma (2016) *Acta Medica Mediterranea*, 32 (1), pp. 209-212. DOI: 10.19193/0393-6384_2016_1_32.
- 73) Özkaya, I., Biçer, E.K., Yardimci, B., Tunçkale, A. A preventive effect of the Mediterranean diet on cardiovascular disease risk in type 2 diabetics (PREMEDIC trial) (2016) *Acta Medica Mediterranea*, 32 (1), pp. 255-261. DOI: 10.19193/0393-6384_2016_1_40.
- 74) Hazavehei, S.M.M., Shahabadi, S., Bashiriyan, S.,

- Karami, M., Almasi, A., Hashemi, S.Z., Saidi, M.R. Profile of cardiovascular diseases risk factors in Kermanshah City, Western Iran: A need assessment study (2016) *Acta Medica Mediterranea*, 32 (Special Issue 2), pp. 995-997.
- 75) Asgary, M.R., Aghajanzadeh, M., Hemmati, H., Jafari, M. Epidemiology of mediastinal tumors during six years (2006-2012) in Rasht City (2016) *Acta Medica Mediterranea*, 32 (Special Issue 4), pp. 1243-1247.
- 76) Shamansouri, S., Karbasizade, V., Khozaie, M. Determining SCCmec types in staphylococcus epidermidis isolated from clinical samples of Ishahan, Iran (2016) *Acta Medica Mediterranea*, 32 (Special Issue 5), pp. 2107-2113.
- 77) Rezaei, S., Velayati, A., Bagheri, A., Moradzadeh, R., Sohrabi, Y., Jamshidi, A., Zare, M., Ansari, G., Ahmadpour, M., Mirzaei, N. Investigation of the levels of air criteria pollutants effecting on health in Dogonbadan City and description of air quality using AQI index (2016) *Acta Medica Mediterranea*, 32 (SpecialIssue4), pp. 1449-1455;.
- 78) Barghamadi, Z., Moosa-Kazemi, S.H., Oshaghi, M.A., Vatandoost, H., Shahraki, G. Species identification of the *Anopheles fluviatilis* complex using phylogenetic analysis PCR-sequencing in Southwestern Iran (2016) *Acta Medica Mediterranea*, 32 (Special Issue 5), pp. 2069-2073.
- 80) Pektas, S.D., Tugbaalatas, E., Yilmaz, N. Plateletcrit is potential biomarker for presence and severity of psoriasis vulgaris (2016) *Acta Medica Mediterranea*, 32 (6), pp. 1785-1790. DOI: 10.19193/0393-6384_2016_6_164.
- 81) Shahtaheri, E., Hosseinian, S., Ebrahimi, M., Jalilabadi, Z., Mirderikvand, F. The impact of stress reduction program based on mindfulness and conscious yoga on depression, perceived stress and quality of life in pregnant women (2016) *Acta Medica Mediterranea*, 32 (SpecialIssue5), pp. 2047-2050.
- 82) Kefeli, A., Yeniova, A.O., Basyigit, S., Uzman, M., Asilturk, Z., Tanrikulu, Y., Nazligül, Y. Effects of helicobacter pylori infections on ghrelin and leptin levels (2016) *Acta Medica Mediterranea*, 32 (5), pp. 1609-1615. DOI: 10.19193/0393-6384_2016_5_139.
- 83) Vardar, E., Basmaci, G., Comut, E., Cavdar, D.K., Erkan, N., Ilhan, E., Vardar, B.U. Comparison of thyroid transcription factor-1 expression in gastric adenocarcinoma and non-cohesive type gastric carcinoma: Is there any difference? (2016) *Acta Medica Mediterranea*, 32 (3), pp. 753-756; DOI: 10.19193/0393-6384_2016_3_85;
- 84) Ulusoy, M.O., Kivanç, S.A., Atakan, M., Uzun, O., Ozturk-Şahin, B., Olcaysu, O.O., Karatepe, A.S., Zorlu-oztürk, S., Turker, I. Are diabetic patients aware of the danger in Turkey? (2016) *Acta Medica Mediterranea*, 32 (4), pp. 925-931. DOI: 10.19193/0393-6384_2016_4_111.
- 85) Güçlü, A., Dursun, B., Rota, S., Sabir, N., Kaya, C., Yaman, F. The relationship between visceral adipose tissue and intima-media thickness in patients with kidney disease (2016) *Acta Medica Mediterranea*, 32 (6), pp. 1843-1849. DOI: 10.19193/0393-6384_2016_6_172.

Corresponding author
MARGHERITA SALERNO, MD
Sciences for Mother and Child Health Promotion
University of Palermo
(Italy)